

Need Money?

Recently Diagnosed?

Desire to help medical research?

Make up to **\$3200 per month** for donating plasma.

■ Hepatitis A

Donor Requirements

- Within three months of initial infection
- Positive test for HAV IgM antibody

Facts about Hepatitis A

- Transmitted by sexual contact, feces contact, or improperly-prepared food
- Symptoms may act flu-like in early stage, muscle ache, headache, appetite loss, abdominal discomfort, fever, jaundice

■ Rubella

German measles

Donor Requirements

- Within three months of initial infection
- Positive test for Rubella IgM antibody

Facts about Rubella

- Transmitted from coughing, sneezing
- Symptoms may include swollen glands, fever, pain or swelling, rash

■ HIV, Syphilis, or Hepatitis B

Donor Requirements

- Within 30 days of initial infection, antigen positive and antibody negative
- Positive test for HIV-2 or 0, or IgM antibody

Facts about HIV, Syphilis and Hepatitis B

- Transmitted by sexual contact, infected blood

**Urgent
Need for
Hep A
Donors**

■ Toxoplasmosis

Donor Requirements

- Within two months of initial infection
- Positive test for Toxo IgM antibody

Facts about Toxoplasmosis

- Transmitted through ingestion of cat feces, contaminated food or drink
- Symptoms may act flu-like in early stage, swollen glands, muscle ache

■ Chagas

Trypanosoma cruzi

Donor Requirements

- Within six months of initial infection
- Positive blood test for Chagas

Facts about Chagas

- Transmitted through blood transfusion or by blood-feeding, triatomine bugs
- Symptoms may include swelling, skin lesion, fever, rapid heartbeat, rash, fatigue, headache, appetite loss

■ Rheumatoid Arthritis

Donor Requirements

- Positive test for Rheumatoid factor greater than 2000 IU/mL

■ Human Parvovirus B19

Donor Requirements

- Within three months of initial infection
- Positive test for HPV antibody

Facts about Human Parvovirus

- Transmitted from coughing, sneezing
- Symptoms may include low-grade fever, rash, nausea, headache, runny nose, skin lesions

■ Mononucleosis

Epstein-Barr virus or cytomegalovirus

Donor Requirements

- Must be within one month of initial infection
- Positive test for EBV IgM or CMV IgM antibody

Facts about Mononucleosis

- Symptoms may include fatigue, appetite loss, chills, sore throat, fever, swollen glands

■ Lyme Disease

Borrelia burgdorferi

Donor Requirements

- Within one year of initial infection
- Positive test for Lyme IgM or Lyme IgG antibody

Donors must be age 18 to 65, weigh at least 110 lbs, and be healthy enough to donate plasma. Translation services available.

Donor Helpline: 888.806.5215
Language interpreters available.

Donor Fax line: 508.244.6593

