

KPL Peroxidase-Labeled Affinity Purified Antibody To Monkey IgG (γ)

Produced in Goat

Cat. No.

5220-0333 (074-11-021)

Size

1.0 mg

DESCRIPTION

Affinity purified antibody isolated from a pool of serum from goats immunized with purified Old World monkey IgG was labeled with horseradish peroxidase using a modified periodate method adapted from Farr and Nakane ⁽¹⁾.

FORM/STORAGE

Lyophilized. Store at 2-8°C until rehydrated. Stable for a minimum of 1 year when stored at 2-8°C.

STABILIZER AND PRESERVATIVE

Bovine serum albumin (BSA) added as a protein stabilizer. A proprietary antibacterial agent is added as a preservative. DO NOT USE SODIUM AZIDE. Non-sterile.

ANTIBODY CONCENTRATION

The concentration of affinity purified antibody is 1 mg as determined by UV absorbance at 280 nm.

E/P RATIO

Molar enzyme/antibody protein ratio = 2:1

SPECIFICITY/CROSS REACTIVITY

Tested by gel diffusion and ELISA techniques as applicable. This product reacts specifically with Old World monkey IgG including Rhesus macaque, chimpanzee, orangutan and Cynomolgus monkey. It has also been found to react strongly with baboon. Product has been adsorbed with purified monkey IgA and IgM. Reactivity to IgG subclasses has not been tested. Antibodies to monkey IgG may cross-react with immunoglobulins of other species if common binding sites are shared.

REHYDRATION STORAGE

Note: Rehydration of antibodies in TBS or buffers other than those listed here is not recommended.

Procedure A: 50% Glycerol

At a working dilution, the level of glycerol is too small to affect most assays. The use of glycerol is not recommended when the conjugate is used in live cell work.

Rehydration: Add 0.5 ml reagent quality water to the product vial. Rotate the vial until the lyophilized pellet is totally dissolved. Add 0.5 ml glycerol to the product vial. Pipet up and down several times to ensure proper mixing. Dilute to desired concentration with TBS or other buffer.

Storage: This product may be stored either refrigerated or frozen as desired. Stable for a minimum of 1 year.

Procedure B: KPL HRP Conjugate Stabilizer

Rehydration: Rehydrate with 1 ml of KPL HRP Conjugate Stabilizer. Rotate the vial until the lyophilized pellet is totally dissolved. Dilute to desired concentration with additional KPL HRP Conjugate Stabilizer.

Storage: This product should be stored at 2-8°C. Stable for a minimum of 1 year.

Procedure C: H₂O

Rehydration: Rehydrate with 1 ml of reagent quality water. Rotate the vial until the lyophilized pellet is totally dissolved. Dilute to desired concentration with TBS or other buffer.

Storage: This product may be stored for up to 1 week refrigerated; thereafter, it should be stored frozen. Stable for a minimum of 1 year at -20°C.

KPL Peroxidase-Labeled Affinity Purified Antibody To Monkey IgG (γ)

Produced in Goat

<u>Cat. No.</u>	<u>Size</u>
5220-0333 (074-11-021)	1.0 mg

SUGGESTED WORKING DILUTIONS

Optimal working concentrations should be determined experimentally. Prepare working dilution in TBS or other buffer such as BSA or KPL Milk Diluent/Blocking Solution (See RELATED PRODUCTS). These buffers not recommended for long term storage. Suggested starting dilutions are as follows. In many cases, the antibody may be diluted further than indicated.

ELISA:

1:1000 – 1:5000 1.0 μ g/ml - 0.2 μ g/ml

Blotting:

1:1000 – 1:5000 1.0 μ g/ml - 0.2 μ g/ml

Histo/Cytochemical Procedures:

1:200 – 1:500 5.0 μ g/ml - 2.0 μ g/ml

PRODUCT SAFETY AND HANDLING

This product is considered non-hazardous as defined by The Hazard Communication Standard (29 CFR 1910.1200). Avoid contact with skin and eyes. In case of contact or spillage, clean with copious amounts of water. Disposal via sanitary sewer.

REFERENCES

1. Farr and Nakane. (1981). J. Immunol. Methods, 47:129-144.

RELATED PRODUCTS

	CAT. NO.
KPL HRP Stabilizer	5290-0005 (54-15-01)
KPL 5X Detector Block	5920-0004 (71-83-00)
KPL Milk Diluent/Blocking Solution	5140-0011 (50-82-01)
KPL Coating Solution	5150-0014 (50-84-00)
KPL Wash Solution	5150-0008 (50-63-00)
KPL SureBlue Microwell Substrate	5120-0075 (52-00-01)
KPL ABTS Microwell Substrate	5120-0041 (50-66-00)
KPL TMB Membrane Substrate	5420-0027 (50-77-03)
KPL HistoMark [®] TrueBlue	5510-0035 (54-78-00)
KPL BSA Diluent/Blocking Solution	5140-0006 (50-61-00)

The product listed herein is for research use only and is not intended for use in human or clinical diagnosis.